

This is a reprint from AAN 2021 virtual congress; the references to “Merck” or “Merck KGaA” within refer to (1) Merck KGaA, Darmstadt, Germany; (2) an affiliate of Merck KGaA, Darmstadt, Germany; or (3) one of the businesses of Merck KGaA, Darmstadt, Germany, which operate as EMD Serono in the healthcare, MilliporeSigma in the life science and EMD Electronics in the electronics business in the U.S. and Canada.

There are two different, unaffiliated companies that use the name “Merck”. Merck KGaA, Darmstadt, Germany, which is providing this reprint for the educational, scientific information contained therein, uses the firm name “Merck KGaA, Darmstadt, Germany” and the business names EMD Serono in the healthcare, MilliporeSigma in the life science and EMD Electronics in the electronics business in the U.S. and Canada. The other company, Merck & Co., Inc. holds the rights in the trademark “Merck” in the U.S. and Canada. Merck & Co., Inc. is not affiliated with or related to Merck KGaA, Darmstadt, Germany, which owns the “Merck” trademark in all other countries of the world.

**GET PRESENTATION PDF**

Copies of this presentation obtained through QR (Quick Response) code are for personal use only and may not be reproduced without written permission of the authors

Characterization of Peripheral Immune Cell Dynamics and Repopulation Patterns in the First 12 Months of Cladribine Tablets Treatment: MAGNIFY-MS Study

H. Wiendl¹, K. Schmierer^{2,3}, S. Hodgkinson⁴, T. Derfuss⁵, A. Chan⁶, F. Sellebjerg⁷, A. Achiron⁸, X. Montalban⁹, A. Prat¹⁰, N. De Stefano¹¹, F. Barkhof^{12,13}, L. Leocani¹⁴, P. Vermersch¹⁵, A. Chudecka¹⁶, S. Roy¹⁷, U. Boschert¹⁷

¹Department of Neurology, Institute of Translational Neurology, University of Münster, Münster, Germany; ²Blizard Institute, Barts and The London School of Medicine & Dentistry, Queen Mary University of London, London, UK; ³Clinical Board Medicine (Neuroscience), The Royal London Hospital, Barts Health NHS Trust, London, UK; ⁴Ingham Institute for Applied Medical Research, University of New South Wales Medicine, Sydney, Australia; ⁵Department of Neurology, University Hospital Basel, Basel, Switzerland; ⁶University Hospital of Berne, University of Berne, Berne, Switzerland; ⁷Danish MS Center, Department of Neurology, University of Copenhagen, Rigshospitalet, Copenhagen, Denmark; ⁸Multiple Sclerosis Center, Sheba Academic Medical Center, Ramat Gan, Israel; ⁹Department of Neurology-Neuroimmunology, Centre d'Esclerosi Múltiple de Catalunya (Cemcat), Hospital Universitari Vall d'Hebron, Barcelona, Spain; ¹⁰Department of Neurosciences, Université de Montréal, Montréal, QC, Canada; ¹¹Department of Neurological and Behavioural Sciences, University of Siena, Siena, Italy; ¹²Department of Radiology, VU University Medical Center, Amsterdam, The Netherlands; ¹³UCL Institute of Neurology, London, UK; ¹⁴Experimental Neurophysiology Unit, Vita-Salute San Raffaele University, Milan, Italy; ¹⁵University of Lille, Inserm U1172 LiNCog, CHU Lille, FHU Precise, Lille, France; ¹⁶Cytel Inc., Geneva, Switzerland; ¹⁷Ares Trading S.A., Eysins, Switzerland, an affiliate of Merck KGaA, Darmstadt, Germany.

Disclosures

This study was sponsored by Merck KGaA, Darmstadt, Germany.

- **HW** is member of Scientific advisory boards/Steering Committees for Bayer, Biogen, Merck KGaA (Darmstadt, Germany), Novartis, Roche, Sanofi-Genzyme, and Teva. He received speaker honoraria and travel support from Bayer, Biogen, CSL Behring, EMD Serono, Fresenius Medical Care, Genzyme, Merck KGaA (Darmstadt, Germany), Omniamed, Novartis, Sanofi-Aventis, and Teva. He received compensation as a consultant from Biogen, Merck KGaA (Darmstadt, Germany), Novartis, Omniamed, Roche, and Sanofi-Genzyme. He has received research support from Bayer, Biogen, Merck KGaA (Darmstadt, Germany), Novartis, Sanofi-Genzyme, Sanofi US, and Teva Pharma, as well as German Ministry for Education and Research (BMBF), German Research Foundation (DFG), Else Kröner Fresenius Foundation, Fresenius Foundation, Hertie Foundation, NRW Ministry of Education and Research, Interdisciplinary Center for Clinical Studies (IZKF) Muenster, and RE Children's Foundation.
- **KS** has been principal investigator of trials sponsored by MedDay, Novartis, Roche, and Teva, and involved in trials sponsored by BIAL, Biogen, Canbex, Cytokinetics, and Genzyme. He has received speaking honoraria from, and/or served in an advisory role for, Biogen, Cinnagen, Merck Inc., Merck KGaA (Darmstadt, Germany), Novartis, Roche, and Teva; and has received research grant support from Biogen and Novartis.
- **SH** serves on advisory boards for Bayer, Biogen, Merck KGaA (Darmstadt, Germany), Novartis, Roche, and Sanofi-Genzyme; she has received money for travel and speaker honoraria from Bayer, Biogen, Merck KGaA (Darmstadt, Germany), Novartis, Roche, and Sanofi-Genzyme.
- **TD** serves on scientific advisory boards for Bayer, Biogen, GeNeuro, MedDay, Merck KGaA (Darmstadt, Germany), Mitsubishi Pharma, Novartis Pharmaceuticals, Roche, and Sanofi-Genzyme; has received funding for travel and/or speaker honoraria from Bayer, Biogen, Merck KGaA (Darmstadt, Germany), Novartis Pharmaceuticals, Roche, and Sanofi-Genzyme; and receives research support from Biogen, the European Union, Novartis Pharma, the Swiss MS Society, and the Swiss National Foundation.
- **ACha** has received speakers'/board honoraria from Actelion (Janssen/J&J), Almirall, Bayer, Biogen, Celgene (BMS), Genzyme, Merck KGaA (Darmstadt, Germany), Novartis, Roche, and Teva, all for hospital research funds; and research support from Biogen, Genzyme, and UCB.
- **FS** has served on scientific advisory boards, been on the steering committees of clinical trials, served as a consultant, received support for congress participation, received speaker honoraria, or received research support for his laboratory from Biogen, EMD Serono, Merck KGaA (Darmstadt, Germany), Novartis, Roche, Sanofi-Genzyme, and Teva.
- **AA** has received honoraria or consulting fees from Bayer, Biogen, Merck KGaA (Darmstadt, Germany), Roche, and Sanofi-Genzyme; and research support from Bayer, Biogen, Merck KGaA (Darmstadt, Germany), Roche, and Sanofi-Genzyme.
- **XM** has received speaking honoraria and travel expenses for participation in scientific meetings, has been a steering committee member of clinical trials or participated in advisory boards of clinical trials in the past years with Actelion (Janssen/J&J), Alexion, Bayer, Biogen, Celgene (BMS), EMD Serono, Genzyme, Immunic, MedDay, Merck KGaA (Darmstadt, Germany), Mylan, Nervgen, Novartis, Roche, Sanofi-Genzyme, Teva Pharmaceutical, TG Therapeutics, Excemed, MSIF and NMSS.
- **AP** has received honoraria and operating grants from pharmaceutical companies.
- **NDS** is a consultant for Bayer, Biogen, Merck KGaA (Darmstadt, Germany), Novartis, Roche, Sanofi-Genzyme, and Teva; has grants or grants pending from FISM and Novartis, is on the speakers' bureaus of Biogen, Merck KGaA (Darmstadt, Germany), Novartis, Roche, Sanofi-Genzyme, and Teva; and has received travel funds from Merck KGaA (Darmstadt, Germany), Novartis, Roche, Sanofi-Genzyme, and Teva.
- **FB** is supported by the NIHR biomedical research centre at UCLH and is a consultant to Roche, Biogen, Merck KGaA (Darmstadt, Germany), IXICO and Combinostics.
- **LL** has received honoraria for consulting services or speaking activities from Biogen, Novartis, Merck KGaA (Darmstadt, Germany), and Roche; and research support from Biogen, Merck KGaA (Darmstadt, Germany), and Novartis.
- **PV** has received honoraria or consulting fees from Almirall, Biogen, Celgene (BMS), MedDay, Merck KGaA (Darmstadt, Germany), Novartis, Roche, Sanofi-Genzyme, and Servier; and research support from Bayer, Biogen, Merck KGaA (Darmstadt, Germany), and Sanofi-Genzyme.
- **AChu** is an employee of Cytel Inc., Geneva Branch, Switzerland, funded by Merck KGaA (Darmstadt, Germany) to perform statistical analyses for this study.
- **SR** and **UB** are employees of Ares Trading S.A., Eysins, Switzerland, an affiliate of Merck KGaA, Darmstadt, Germany.

Medical writing assistance was provided by Joe Ward of inScience Communications, Springer Healthcare Ltd, UK, and was funded by Merck KGaA, Darmstadt, Germany.

The MAGNIFY-MS study: NCT03364036.

Introduction

- In the treatment of MS, immune reconstitution therapies are used for short, intermittent periods to allow for treatment-free periods.^{1,2}
- Long-term lymphocyte dynamics have been evaluated for CLARITY, CLARITY Extension, and PREMIERE studies,³ indicating immune cell repopulation after treatment with cladribine tablets.
- MAGNIFY-MS aims to determine the onset of action of cladribine tablets (3.5 mg/kg cumulative dose over 2 years) in patients with highly active relapsing MS[†].
 - The action of cladribine tablets on immune cells may be key for both onset and durability of its effect in people with MS.

[†]Highly active relapsing MS as defined by: one relapse in the previous year and at least one T1 Gd+ lesion, or nine or more T2 lesions, while on therapy with other DMTs, or two or more relapses in the previous year, whether on DMT treatment or not.

1. Wiendl H. *Nat Rev Neurol*. 2017;13:573–574. 2. Giovannoni G. *Neurotherapeutics*. 2017;14:874–887. 3. Comi G, et al. *Mult Scler Relat Disord*. 2019;29:168–174.
DMT, disease-modifying therapy; Gd+, gadolinium enhancing; MRI, magnetic resonance imaging; MS, multiple sclerosis

Introduction

- In the treatment of MS, immune reconstitution therapies are used for short, intermittent periods to allow for treatment-free periods.^{1,2}
- Long-term lymphocyte dynamics have been evaluated for CLARITY, CLARITY Extension, and PREMIERE studies,³ indicating immune cell repopulation after treatment with cladribine tablets.


OBJECTIVE

To report on peripheral immune cell subset dynamics and immunoglobulin levels in the first 12 months of cladribine tablets therapy.

¹Highly active relapsing MS as defined by: one relapse in the previous year and at least one T1 Gd+ lesion, or nine or more T2 lesions, while on therapy with other DMTs, or two or more relapses in the previous year, whether on DMT treatment or not.

1. Wiendl H. *Nat Rev Neurol*. 2017;13:573–574. 2. Giovannoni G. *Neurotherapeutics*. 2017;14:874–887. 3. Comi G, et al. *Mult Scler Relat Disord*. 2019;29:168–174.
DMT, disease-modifying therapy; Gd+, gadolinium enhancing; MRI, magnetic resonance imaging; MS, multiple sclerosis


Methods



MAGNIFY-MS is an ongoing Phase IV, open-label, single-arm, multicenter, 2-year study. Patients with highly active relapsing MS† received cladribine tablets, with 2 weeks active treatment per course (Week 1 and Week 5 of each year).

†Highly active relapsing MS as defined by: one relapse in the previous year and at least one T1 Gd+ lesion, or nine or more T2 lesions, while on therapy with other DMTs, or two or more relapses in the previous year, whether on DMT treatment or not.
9HPT, 9-hole peg test; **BL**, baseline; **CUA**, combined unique active; **DMT**, disease-modifying therapy; **EDSS**, Expanded Disability Status Scale; **HIV**, human immune deficiency virus; **KFS**, Kurtzke Functional System; **MRI**, magnetic resonance imaging; **MS**, multiple sclerosis; **SDMT**, symbol digit modalities test; **T25FW**, timed 25-foot walk; **TB**, tuberculosis

Methods



- This sub-study of MAGNIFY-MS involved longitudinal evaluation of peripheral blood immune cells in patients receiving cladribine tablets.
- 57 patients were treated.
- Absolute cell counts and % change from baseline were assessed for adaptive immune cell subtypes and immunoglobulins.
- Immunophenotyping was completed at baseline and at months 1*, 2*, 3, 6, and 12.

*Months 1 and 2 are available for TBNK and B cell panels.

BL, baseline; Breg, B regulatory; Ig, immunoglobulin; TEMRA, terminally differentiated effector memory RA⁺; Treg, T regulatory

Methods


Immune cell panels in MAGNIFY-MS

T cell panel	<p>CD4 naïve (CD3⁺, CD8⁻, CD4⁺, CD45RA⁺, CCR7⁺)</p> <p>CD4 central memory (CD3⁺, CD8⁻, CD4⁺, CD45RA⁻, CCR7⁺)</p> <p>CD4 effector memory (CD3⁺, CD8⁻, CD4⁺, CD45RA⁻, CCR7⁻)</p> <p>CD8 naïve (CD3⁺, CD4⁻, CD8⁺, CD45RA⁺, CCR7⁺)</p> <p>CD8 central memory (CD3⁺, CD4⁻, CD8⁺, CD45RA⁻, CCR7⁺)</p> <p>CD8 effector memory (CD3⁺, CD4⁻, CD8⁺, CD45RA⁻, CCR7⁻)</p> <p>CD8 TEMRA (CD3⁺, CD4⁻, CD8⁺, CD45RA⁺, CCR7⁻)</p> <p>Th1 (CD3⁺, CD8⁻, CD4⁺, CCR7^{-/+}, CXCR3⁺)</p> <p>Treg (CD3⁺, CD8⁻, CD4⁺, CD25^{bright}, CD127^{dim/-})</p> <p>Th17 (CD3⁺, CD8⁻, CD4⁺, CD45RA⁻, CCR7^{-dim}, CCR6⁺, CD146⁺)</p> <p>Th2 (CD3⁺, CD8⁻, CD4⁺, CXCR3⁺, CCR6⁻)</p>	B cell panel	<p>CD19 B cells (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD19⁺)</p> <p>CD20 B cells (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD20⁺)</p> <p>Activated B cells (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD19⁺, CD20⁺, CD69⁺)</p> <p>Naïve B cells (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD19⁺, CD20⁺, IgD⁺, CD27⁻)</p> <p>Memory B cells (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD19⁺, CD20⁺, CD27⁺)</p> <p>Short-lived plasma cells (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD19^{dim}, CD20^{-/dim}, CD27^{bright})</p> <p>Breg (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD19⁺, CD24^{bright}, CD38^{bright})</p> <p>CD38^{bright} plasma cells (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD19^{dim}, CD20⁻, CD38^{bright})</p> <p>Transitional B cells (CD45^{bright}, SSC^{low}, CD3⁻, CD14⁻, CD56⁻, CD19⁺, CD20⁺, IgD⁺, CD10⁺, CD27⁻)</p>
NK cell panel	<p>NKp46 NK cells (SSC^{low}, CD45⁺, CD19⁻, CD3⁻, CD16⁺ CD56⁺, CD335⁺)</p> <p>CD16 NK cells (SSC^{low}, CD45⁺, CD19⁻, CD3⁻, CD16⁺ CD56⁺)</p> <p>CD16^{bright} CD56^{dim} NK cells (SSC^{low}, CD45⁺, CD19⁻, CD3⁻, CD56^{dim}, CD16^{bright})</p> <p>CD16^{low} CD56^{bright} NK cells (SSC^{low}, CD45⁺, CD19⁻, CD3⁻, CD56^{bright}, CD16^{-/+})</p> <p>CD16⁻CD56^{dim} NK cells (SSC^{low}, CD45⁺, CD19⁻, CD3⁻, CD56^{dim}, CD16⁻)</p> <p>CD16⁺ CD56⁻ NK cells (SSC^{low}, CD45⁺, CD19⁻, CD3⁻, CD56⁻, CD16⁺)</p> <p>CD16⁺ CD56⁺ NK cells</p>		
Immunoglobulins	<p>IgG</p> <p>IgM</p>		


Results

Absolute values ($10^6/L$) of B and T cells





B cell subtypes



CD4⁺ T cell subtypes



CD8⁺ T cell subtypes




- CD19⁺ B cells
- CD20⁺ B cells
- Activated B cells (CD69⁺)
- Naïve B cells
- Memory B cells
- Short-lived plasma cells
- Regulatory B cells
- CD38⁺ plasma cells
- Transitional B cells

- CD4⁺ T cells
- Central memory T cells
- Effector memory T cells
- Naïve T cells
- Regulatory T cells
- Th1 cells
- Th17 cells
- Th2 cells

- CD8⁺ T cells
- Central memory T cells
- Effector memory T cells
- Naïve T cells
- TEMRA T cells



Results

Absolute values (g/L) of immunoglobulins



Results

Absolute values ($10^6/L$) of Natural Killer cells



- CD16⁺ CD56⁺
- CD16^{bright} CD56^{dim}
- CD16^{low} CD56^{bright}
- CD16
- CD16⁺ CD56⁻
- NKp46 cells
- CD16⁻ CD15^{dim}

Results

Key:

Nadir value

Values above
baseline level

Percentage change from baseline of B cell subtypes

Subtype	Month 1	Month 2	Month 3	Month 6	Month 12
CD19 ⁺ B cells	-77% n=46	-90% n=44	-80% n=46	-60% n=35	-35% n=42
Memory B cells	-74% n=45	-93% n=44	-93% n=46	-90% n=34	-87% n=42
Activated B cells (CD69 ⁺)	-64% n=45	-81% n=44	-73% n=46	-53% n=34	-45% n=42
CD38 ⁺ plasma cells	-11% n=45	-66% n=44	-71% n=46	-51% n=34	-51% n=42
Short-lived plasma cells	-28% n=45	-65% n=44	-78% n=46	-58% n=34	-51% n=42
Naïve B cells	-80% n=45	-90% n=44	-75% n=46	-43% n=34	-5% n=42
Transitional B cells	-61% n=45	-63% N=44	+28% n=46	+34% n=34	+36% n=42
Regulatory B cells	-45% n=45	-16% n=44	+176% n=46	+171% n=34	+50% n=42

- There is an early onset of action, with most B cell subtypes reaching nadir levels by month 2.
- Reduction in memory B cells was sustained to month 12; regulatory B cells recovered by month 3, and then increased over baseline levels.

Results

Key:

Nadir value

Values above
baseline level

Percentage change from baseline of T cell subtypes

Subtype	Month 1	Month 2	Month 3	Month 6	Month 12
CD4+	-22% n=46	-51% n=44	-54% n=46	-51% n=35	-40% n=42
CD8+	-18% n=46	-39% n=44	-50% n=46	-43% n=35	-36% n=42
CD4+ Th1	-	-	-51% n=46	-45% n=34	-35% n=42
CD4+ Th17	-	-	-35% n=45	-34% n=34	-24% n=41
CD4+ CM/EffM	-	-	-56%/-35% n=46	-52%/-35% n=34	-40%/-15% n=42
CD8+ CM/EffM	-	-	-39%/-29% n=46	-46%/-22% n=34	-29%/-16% n=42
CD4+ naïve/CD8+ naïve	-	-	-63%/-73% n=46/n=46	-64%/-70% n=35/n=34	-53%/-58% n=42/n=42
CD4+ Treg	-	-	-30% n=46	-38% n=34	-21% n=42
CD8+ TEMRA	-	-	-26% n=46	-15% n=34	-19% n=42

- T cell subtypes show reductions at later time points.

Results

Key:

Nadir value

Values above
baseline level

Percentage change from baseline of NK cell subtypes

Subtype	Month 1	Month 2	Month 3	Month 6	Month 12
CD16 ^{low} CD56 ^{bright}	-	-	-7% n=46	-2% n=34	-9% n=42
CD16 ⁺ CD56 ⁺	-34% n=46	-40% n=44	-34% n=46	-17% n=35	-14% n=42
CD16 ^{bright} CD56 ^{dim}	-	-	-40% n=46	-19% n=34	-12% n=42

- No changes were seen for other NK cell subtypes.

Conclusions

- **MAGNIFY-MS MRI results suggest an early onset of cladribine tablets action. This may be mediated through a specific pattern of sustained decrease and reconstitution of B and T cell subtypes.**
- **The pronounced effect on B cells, especially memory B cells in the first 2 months of cladribine tablet treatment, suggests a contribution to early efficacy onset.**
- **Sustained depletion of memory B cells and the moderate decrease across T cell subtypes may contribute to the long-term effect of cladribine tablets.**