

“This reprint might contain references to “Merck” or “Merck KGaA”, which refer to (1) Merck KGaA, Darmstadt, Germany; (2) an affiliate of Merck KGaA, Darmstadt, Germany; or (3) one of the businesses of Merck KGaA, Darmstadt, Germany, which operate as EMD Serono in the healthcare, MilliporeSigma in the life science and EMD Electronics in the electronics business in the U.S. and Canada.

There are two different, unaffiliated companies that use the name “Merck”. Merck KGaA, Darmstadt, Germany, which is providing this content, uses the firm name “Merck KGaA, Darmstadt, Germany” and the business names EMD Serono in the healthcare, MilliporeSigma in the life science and EMD Electronics in the electronics business in the U.S. and Canada. The other company, Merck & Co., Inc. holds the rights in the trademark “Merck” in the U.S. and Canada. Merck & Co., Inc. is not affiliated with or related to Merck KGaA, Darmstadt, Germany, which owns the “Merck” trademark in all other countries of the world.”

Long-term Disease Stability Assessed by the Expanded Disability Status Scale in Patients Treated with Cladribine Tablets in the CLARITY and CLARITY Extension Studies

G. Giovannoni,¹ G. Comi,² K. Rammohan,³ P. Rieckmann,⁴ P. Vermersch,⁵ F. Dangond,⁶ B. Keller,⁷ D. Jack⁷

¹Queen Mary University of London, Blizard Institute, Barts and The London School of Medicine and Dentistry, London, UK; ²Department of Neurology and Institute of Experimental Neurology, Università Vita-Salute San Raffaele, Ospedale San Raffaele, Milan, Italy; ³University of Miami School of Medicine, Department of Neurology, MS Research Center, Clinical Research Building, Miami, FL, USA; ⁴Department of Neurology, Medical Park Loipl, and University of Erlangen, Erlangen, Germany; ⁵Univ. Lille, INSERM U995, CHU Lille, FHU Imminent, Lille, France; ⁶EMD Serono Research & Development Institute, Inc., Billerica, MA, USA, an affiliate of Merck KGaA, Darmstadt, Germany; ⁷Merck KGaA, Darmstadt, Germany



GET POSTER PDF
Copies of this poster obtained through QR (Quick Response) code are for personal use only and may not be reproduced without written permission of the authors.




RESULTS (cont.)

Annual EDSS Stability

- In the CP3.5 group, in each yearly period, EDSS score was stable in 53.9–76.5% of patients (**Figure 3**)
- In each yearly period, EDSS score improved in 21–30% of patients and worsened in 0–25%
- During Year 5, EDSS score stability was observed in 48 (53.9%) patients, improvement in 19 (21.3%), and worsening in 22 (24.7%) in the CP3.5 group (n=89)
- In the CC7 group during Year 5 (n=171), EDSS score remained stable in 113 patients (66.1%), improved in 31 (18.1%), and worsened in 27 (15.8%)

Figure 3. Change in EDSS Score in Each 12 Month Period Up to 5 Years in the CP3.5 and CC7 Patient Groups




Improved: any decrease in EDSS score ≥ 1 (baseline EDSS 1–4.5) OR ≥ 0.5 (EDSS ≥ 5.0).
Stable: all remaining cases i.e. increase or decrease < 1.0 (baseline EDSS 0.5–4.5) OR < 1.5 (EDSS 0) OR no change (EDSS ≥ 5.0).
Worsened: any increase in EDSS score ≥ 1 (baseline EDSS 0.5–4.5) OR ≥ 1.5 (EDSS 0) OR ≥ 0.5 (EDSS ≥ 5.0).

Time to EDSS Progression

- In both the CP3.5 and CC7 groups, less than 30% of subjects reached 3-month confirmed EDSS progression by 5 years (**Figure 4A**)
- A similar pattern was observed with time to 6-month confirmed EDSS progression. Less than 30% of subjects had confirmed EDSS progression in both the CP3.5 and CC7 groups (**Figure 4B**)

Figure 4. Time to 3- (A) and 6-Month (B) Confirmed EDSS Progression



CONCLUSIONS



Median EDSS score remained stable for up to 5 years post-CLARITY baseline in both the CP3.5 and CC7 treatment groups



Over 50% of patients had stable EDSS score up to 5 years from baseline



Less than 30% of subjects reached 3- or 6-month confirmed EDSS progression at 5 years

INTRODUCTION

- The CLARITY study demonstrated that treatment with cladribine tablets 10 mg (cumulative dose 3.5 mg/kg over 2 years, henceforth referred to as cladribine tablets 3.5 mg/kg [CT3.5]) significantly reduced relapse rates and slowed disability progression versus placebo in RRMS patients¹
- Moreover, the CLARITY Extension study concluded that treatment with cladribine tablets for 2 years followed by treatment with placebo for 2 years produced similar clinical benefits to 4 years of cladribine tablets treatment but with lower incidence of grade 3/4 lymphopenia²
- Disease stability in MS can be assessed through the EDSS. The scale for neurological impairment ranges from 0–10 where a higher score indicates a greater degree of disability and therefore progression of the disease³

OBJECTIVE


- To evaluate *post hoc*, long-term disease stability assessed by the EDSS score after treatment with CT3.5 in patients with RRMS enrolled in CLARITY and CLARITY Extension

METHODS

Eligibility and Endpoints

- Patients enrolled into CLARITY Extension who were randomized to CT3.5 in CLARITY with at least one post-baseline EDSS measurement were included for analysis
 - CLARITY Extension was not a pre-planned study; after completing CLARITY there was a variable bridging interval (median duration: 43 weeks) between studies where no cladribine tablets were administered²
 - Two treatment groups were investigated (**Figure 1**):
 - CP3.5: CT3.5 in CLARITY followed by placebo in CLARITY Extension
 - CC7: CT3.5 in CLARITY followed by CT3.5 in CLARITY Extension
- Endpoints:
 - EDSS scores over time at 6-monthly intervals, from CLARITY randomization to end of follow-up in CLARITY Extension, including the interval between studies
 - Many patients whose EDSS score is shown for CLARITY are not represented in the values of CLARITY Extension
 - No EDSS scores were retrospectively collected during the bridging interval between CLARITY and CLARITY Extension
 - Time to 3- and 6-month confirmed EDSS progression from CLARITY randomization
- EDSS score improvement or worsening each year was defined as any increase or decrease in minimum EDSS score at 6-monthly intervals. All other cases were classified as stable
- An increase or decrease was defined as EDSS score changes of:
 - At least 1.5 points, if baseline EDSS score = 0
 - At least 1 point, if baseline EDSS score ≤ 4.5
 - At least 0.5 points, if baseline EDSS score ≥ 5

Figure 1. CLARITY/CLARITY Extension Study Arms Under Analysis



The only approved dose of cladribine tablets is 3.5 mg/kg of body weight over 2 years. Following completion of the two treatment courses, no further cladribine treatment is required in Years 3 and 4.

METHODS (cont.)

Statistical Analyses


- This was a *post hoc* exploratory analysis of a subset of patients who completed 6 years of follow-up
- EDSS score over time was descriptively analysed (median, 95% CI at 6-monthly intervals, overall and by bridging interval length (≤ 43 weeks versus > 43 weeks))
- Improvement, worsening and stability of EDSS score over 12 months were descriptively analysed (n, %)
- Three and 6-month EDSS progression from CLARITY entry was analyzed by Kaplan-Meier plots

RESULTS

EDSS over Time

- Five years after CLARITY baseline, including variable bridging interval, median EDSS score remained stable compared with baseline values for both groups (**Figure 2**)
 - Median EDSS score remained between 2.0–3.0 up to 60 months in the CP3.5 group (n=69). Median change in EDSS score was 0 points up to 60 months
 - In the CC7 group (n=126), median EDSS score ranged between 2.0–2.5 up to 60 months
- Median (95% CI) EDSS score for patients in the CP3.5 group at 5 years was 2.5 (2.0–3.5) compared with 3.0 (2.5–3.5) at baseline
- In the CC7 group, median EDSS score (95% CI) was 2.0 (2.0–3.0) compared with 2.5 (2.5–3.0) at baseline
- Similar results were observed for both groups when split by variable bridging interval of \leq or $>$ 43 weeks

Figure 2. EDSS Score Over Time in Patients Treated with CP3.5 and CC7



Line = Median. Circles = Mean. Box = Q1, Q3. Error bars reflect the minimum and maximum EDSS score.

Abbreviations: CC7, CT3.5 CLARITY, CT3.5 CLARITY Extension; CP3.5, CT3.5 CLARITY, placebo CLARITY Extension; CI, confidence interval; D1, randomization date; EDSS, Expanded Disability Status Scale; MS, multiple sclerosis; RRMS, relapsing-remitting multiple sclerosis

Acknowledgments: This study was sponsored by EMD Serono, Inc., USA, an affiliate of Merck KGaA, Darmstadt, Germany. Medical writing assistance was provided by Matthew Bexon and Joseph Ward of InScience Communications, Springer Healthcare, Chester, UK and funded by Merck KGaA, Darmstadt, Germany.

Disclosures: GG has received speaker honoraria and consulting fees from Abbvie, Actelion, Atara Bio, Almirall, Bayer Schering Pharma, Biogen Idec, FivePrime, GlaxoSmithKline, GW Pharma, Merck & Co., Merck KGaA (Darmstadt, Germany), Pfizer Inc, Protein Discovery Laboratories, Teva Pharmaceutical Industries Ltd, Sanofi-Genzyme, UCB, Vertex Pharmaceuticals, Ironwood, and Novartis; and has received research support unrelated to this study from Biogen Idec, Merck & Co., Novartis, and Ironwood. GC has received consulting fees from Novartis, Teva Pharmaceutical Industries Ltd., Sanofi-Aventis, Merck KGaA (Darmstadt, Germany), Receptos, Biogen Idec, Genentech-Roche, and Bayer Schering; lecture fees from Novartis, Teva Pharmaceutical Ind. Ltd., Sanofi-Aventis, Merck KGaA (Darmstadt, Germany), Biogen Dompé, Bayer Schering, and Serono Symposia International Foundation; and trial grant support from Novartis, Teva Pharmaceutical Ind. Ltd., Sanofi-Aventis, Receptos, Biogen Idec, Genentech-Roche, Merck KGaA (Darmstadt, Germany), Biogen Dompé, Bayer Schering, KR has received honoraria for lectures/steering committee meetings from EMD Serono, Inc., USA (an affiliate of Merck KGaA, Darmstadt, Germany), Biogen Idec, Sanofi-Aventis, Genzyme, Novartis, Teva Pharmaceutical Industries, and Serono Symposia International Foundation. PR has received honoraria for lectures/steering committee meetings from Merck KGaA (Darmstadt, Germany), Biogen Idec, Bayer Schering Pharma, Boehringer-Ingelheim, Sanofi-Aventis, Genzyme, Novartis, Teva Pharmaceutical Industries, and Serono Symposia International Foundation. PV has received honoraria or consulting fees from Biogen, Sanofi-Genzyme, Bayer, Novartis, Merck KGaA (Darmstadt, Germany), Celgene, Roche and Almirall; and research support from Biogen, Sanofi-Genzyme, Bayer, and Merck KGaA (Darmstadt, Germany). FD is an employee of EMD Serono Research & Development Institute, Inc., USA, an affiliate of Merck KGaA, Darmstadt, Germany. BK and DJ are employees of Merck KGaA, Darmstadt, Germany.

The CLARITY study: NCT00213135. The CLARITY Extension study: NCT00641537.

References: 1. Giovannoni G, et al. *N Engl J Med*. 2010;362:416–426. 2. Giovannoni G, et al. *Mult Scler J*. 2018;24:1594–1604.

Presented at the CMSC 2020 Virtual Poster/Platform Session, August 3, 2020