

# Comparative Assessment of Switching Outcomes for Patients With Multiple Sclerosis Receiving Cladribine Tablets vs Fingolimod, Dimethyl fumarate, and Teriflunomide

Carroline Lobo,<sup>1</sup> Xiaoxue Chen,<sup>1</sup> Xin Zhao,<sup>2</sup> James Cutler,<sup>2</sup> Jeffrey Anderson,<sup>2</sup> Sonia Kim,<sup>2</sup> Emily Evans,<sup>1</sup> Amy L. Phillips,<sup>1</sup> Gabriel Gomez Rey<sup>2</sup>

<sup>1</sup>EMD Serono, Rockland, MA, USA; <sup>2</sup>Genesis Research Group, Hoboken, NJ, USA

## CONCLUSIONS

- In this real-world study, fewer patients receiving cladribine tablets switched to another disease-modifying therapy (DMT) over 2 years compared with patients receiving fingolimod, dimethyl fumarate, or teriflunomide**
- Odds of switching and hazard for time to switching were significantly greater with fingolimod, dimethyl fumarate, and teriflunomide vs cladribine tablets**

## BACKGROUND

- Cladribine tablets are the first and only FDA-approved treatment for relapsing remitting multiple sclerosis (RRMS) and active secondary progressive multiple sclerosis (SPMS) that provides 2 years of proven efficacy with a maximum of 20 days of oral treatment over 96 weeks
- Comparative data for cladribine tablets vs other oral disease-modifying therapies (DMTs) in patients with multiple sclerosis (MS) are emerging

## OBJECTIVE

- To evaluate treatment switching within 1- and 2-year follow-up in patients in the United States with MS treated with cladribine tablets vs fingolimod, dimethyl fumarate, and teriflunomide

## METHODS

- This administrative claims study used data from the Komodo Healthcare Map™ database
- Inclusion criteria were: ≥2 MS diagnoses ≥30 days apart between 4/1/2018 and 12/31/2022; ≥1 claim for cladribine tablets, fingolimod, dimethyl fumarate, or teriflunomide between 4/1/2019 and 12/31/2020 (initiation date = index); no index DMT 1 year before index (baseline); age ≥18 years; and continuous enrollment with healthcare insurance from baseline to 2 years after index (follow-up)
- In each comparison, propensity score weighting was used to balance covariates, including age, sex, index year, race/ethnicity, geographic region, insurance, Charlson Comorbidity Index (CCI) score, MS severity, baseline relapses, MS-related hospitalizations, baseline medications, and MS-related comorbidities
- Weighting by odds ratio (OR) was used to estimate the average treatment effect among those treated with cladribine tablets
- The proportions of patients switching to another DMT within 1 and 2 years (weighted ORs with 95% CIs) and time to switch (weighted proportional hazards Cox regression model hazard ratios [HRs]) were compared across DMTs. Covariates remaining unbalanced after weighting were included in the models

# RESULTS (cont.)

- The overall mean (SD) age was 45 (11) years and 75.7% of patients were female. Most patients had commercial health insurance (63.6%), followed by Medicaid (26.7%), and Medicare (9.7%) (Table 1)

### Table 1. Patient Baseline Demographic Characteristics

|                                       | Overall     | Weighted Population |            |                   |               |
|---------------------------------------|-------------|---------------------|------------|-------------------|---------------|
|                                       |             | Cladribine tablets  | Fingolimod | Dimethyl fumarate | Teriflunomide |
| Total, N                              | 4709        | 269                 | 264        | 253               | 255           |
| <b>Age at index (years)</b>           |             |                     |            |                   |               |
| Mean (SD)                             | 45 (11)     | 44 (11)             | 43 (12)    | 44 (12)           | 44 (11)       |
| <b>Sex, N (%)</b>                     |             |                     |            |                   |               |
| Female                                | 3564 (75.7) | 209 (77.7)          | 208 (78.8) | 194 (76.7)        | 197 (77.3)    |
| Male                                  | 1145 (24.3) | 60 (22.3)           | 56 (21.2)  | 59 (23.3)         | 58 (22.7)     |
| <b>Index year, N (%)</b>              |             |                     |            |                   |               |
| 2019                                  | 2447 (52.0) | 86 (32.0)           | 97 (36.7)  | 83 (32.8)         | 83 (32.5)     |
| 2020                                  | 2262 (48.0) | 183 (68.0)          | 167 (63.3) | 170 (67.2)        | 172 (67.5)    |
| <b>Race/ethnicity, N (%)</b>          |             |                     |            |                   |               |
| White                                 | 2013 (42.7) | 114 (42.4)          | 103 (39.0) | 116 (45.8)        | 125 (49.0)    |
| Black or African American             | 565 (12.0)  | 32 (11.9)           | 40 (15.2)  | 31 (12.3)         | 22 (8.6)      |
| Asian or Pacific Islander             | 45 (1.0)    | 3 (1.1)             | 2 (0.8)    | 2 (0.8)           | 1 (0.4)       |
| Hispanic or Latino                    | 383 (8.1)   | 31 (11.5)           | 29 (11.0)  | 18 (7.1)          | 24 (9.4)      |
| Other                                 | 164 (3.5)   | 10 (3.7)            | 8 (3.0)    | 9 (3.6)           | 8 (3.1)       |
| Unknown                               | 1539 (32.7) | 79 (29.4)           | 81 (30.7)  | 77 (30.4)         | 75 (29.4)     |
| <b>Geographic region, N (%)</b>       |             |                     |            |                   |               |
| Midwest                               | 1150 (24.4) | 60 (22.3)           | 52 (19.7)  | 53 (20.9)         | 53 (20.8)     |
| Northeast                             | 1179 (25.0) | 70 (26.0)           | 61 (23.1)  | 67 (26.5)         | 71 (27.8)     |
| South                                 | 1581 (33.6) | 102 (37.9)          | 107 (40.5) | 97 (38.3)         | 98 (38.4)     |
| West                                  | 799 (17.0)  | 37 (13.8)           | 44 (16.7)  | 36 (14.2)         | 34 (13.3)     |
| <b>Insurance type at index, N (%)</b> |             |                     |            |                   |               |
| Commercial                            | 2994 (63.6) | 171 (63.6)          | 161 (61.0) | 162 (64.0)        | 164 (64.3)    |
| Medicare                              | 456 (9.7)   | 32 (11.9)           | 32 (12.1)  | 31 (12.3)         | 32 (12.5)     |
| Medicaid                              | 1259 (26.7) | 66 (24.5)           | 71 (26.9)  | 61 (24.1)         | 59 (23.1)     |

- Mean (SD) CCI score was 0.55 (1.11) (Table 2)

### Table 2. Patient Baseline Clinical Characteristics

|  | Overall     | Weighted Population |             |                   |               |
|--|-------------|---------------------|-------------|-------------------|---------------|
|  |             | Cladribine tablets  | Fingolimod  | Dimethyl fumarate | Teriflunomide |
| Total, N                                 | 4709        | 269                 | 264         | 253               | 255           |
| <b>Charlson Comorbidity Index score</b>  |             |                     |             |                   |               |
| Mean (SD)                                | 0.55 (1.11) | 0.76 (1.33)         | 0.87 (1.75) | 0.75 (1.34)       | 0.75 (1.41)   |
| <b>Baseline MS severity, N (%)</b>       |             |                     |             |                   |               |
| MS relapse                               | 1664 (35.3) | 115 (42.8)          | 126 (47.7)  | 106 (41.9)        | 110 (43.1)    |
| MS hospitalization                       | 354 (7.5)   | 24 (8.9)            | 34 (12.9)   | 23 (9.1)          | 20 (7.8)      |
| <b>Baseline MS Severity Score</b>        |             |                     |             |                   |               |
| Mean (SD)                                | 4.3 (3.6)   | 5.7 (4.2)           | 6.1 (5.0)   | 5.4 (3.8)         | 5.5 (4.1)     |
| <b>Baseline MS Severity Score, N (%)</b> |             |                     |             |                   |               |
| Low (0-4)                                | 2832 (60.1) | 127 (47.2)          | 129 (48.9)  | 117 (46.2)        | 123 (48.2)    |
| Moderate (5-10)                          | 1554 (33.3) | 100 (37.2)          | 85 (32.2)   | 108 (42.7)        | 94 (36.9)     |
| High (≥11)                               | 323 (6.9)   | 42 (15.6)           | 50 (18.9)   | 28 (11.1)         | 38 (14.9)     |
| <b>Daifampridine use, N (%)</b>          | 141 (3.0)   | 32 (11.9)           | 30 (11.4)   | 24 (9.5)          | 22 (8.6)      |
| <b>Overall DMT use, N (%)</b>            | 1653 (35.1) | 134 (49.8)          | 135 (51.1)  | 118 (46.6)        | 120 (47.1)    |
| <b>DMT use by drug class, N (%)</b>      |             |                     |             |                   |               |
| Oral                                     | 139 (3.0)   | 17 (6.3)            | 12 (4.5)    | 4 (1.6)           | 9 (3.5)       |
| Monoclonal antibody                      | 222 (4.7)   | 61 (22.7)           | 62 (23.5)   | 47 (18.6)         | 51 (20.0)     |
| Platform                                 | 1345 (28.6) | 61 (22.7)           | 65 (24.6)   | 75 (29.6)         | 69 (27.1)     |

DMT, disease-modifying therapy; MS, multiple sclerosis.

- For the weighted population, proportions of patients switching within 1 year were 3.7% on cladribine tablets, 17.8% on fingolimod, 28.9% on dimethyl fumarate, and 21.2% on teriflunomide (Table 3)
- The ORs (95% CI) of 1-year switch vs cladribine tablets were fingolimod 5.66 (2.90-12.15), dimethyl fumarate 10.81 (5.66-22.87), and teriflunomide 7.34 (3.78-15.74) (Table 4)
- The HRs (95% CIs) for time to switching at 1 year were statistically significant compared with cladribine tablets (fingolimod 5.21 [2.54-10.67], dimethyl fumarate 9.26 [4.91-17.45], and teriflunomide 6.65 [3.46-12.79]) (Table 4)

## LIMITATIONS

- Limitations of this study include the relatively small sample size for patients receiving cladribine tablets and the lack of direct clinical measures of MS disease severity and relapse
- The observed results may not be generalizable to other patient populations due to potential differences in data source populations, indications, treatment practices, and endpoint definitions

### Table 3. Switching Outcomes Within 1 Year

|                                       | Cladribine tablets N (%) | Fingolimod N (%) | Dimethyl fumarate N (%) | Teriflunomide N (%) |
|---------------------------------------|--------------------------|------------------|-------------------------|---------------------|
| Total, N                              | 269                      | 264              | 253                     | 255                 |
| <b>Switch within 1 year</b>           | 10 (3.7)                 | 47 (17.8)        | 73 (28.9)               | 54 (21.2)           |
| Total number of patients who switched | 10                       | 47               | 73                      | 54                  |
| <b>Name of DMT switched to</b>        |                          |                  |                         |                     |
| Platform                              | <b>2 (0.7)</b>           | <b>7 (2.6)</b>   | <b>14 (5.5)</b>         | <b>13 (5.1)</b>     |
| Glatiramer                            | 0 (0.0)                  | 5 (2.0)          | 9 (3.4)                 | 8 (3.1)             |
| Interferon beta-1a                    | 2 (0.7)                  | 0 (0.0)          | 3 (1.4)                 | 4 (1.7)             |
| Interferon beta-1b                    | 0 (0.0)                  | 0 (0.0)          | 1 (0.2)                 | 1 (0.3)             |
| Peginterferon beta-1b                 | 0 (0.0)                  | 1 (0.3)          | 1 (0.5)                 | 0 (0.0)             |
| Monoclonal antibodies                 | <b>6 (2.2)</b>           | <b>23 (8.7)</b>  | <b>28 (11.0)</b>        | <b>23 (9.0)</b>     |
| Alemtuzumab                           | 1 (0.4)                  | 0 (0.0)          | 0 (0.0)                 | 0 (0.0)             |
| Natalizumab                           | 0 (0.0)                  | 3 (1.0)          | 7 (2.9)                 | 7 (2.7)             |
| Ocrelizumab                           | 2 (0.7)                  | 20 (7.5)         | 19 (7.5)                | 14 (5.7)            |
| Oratumumab                            | 3 (1.1)                  | 1 (0.2)          | 1 (0.5)                 | 2 (0.6)             |
| Oral                                  | <b>2 (0.7)</b>           | <b>17 (6.5)</b>  | <b>31 (12.0)</b>        | <b>18 (7.1)</b>     |
| Cladribine tablets                    | 0 (0.0)                  | 2 (0.6)          | 1 (0.4)                 | 2 (0.9)             |
| Dimethyl fumarate                     | 1 (0.4)                  | 7 (2.8)          | 0 (0.0)                 | 7 (2.9)             |
| Diroximel fumarate                    | 0 (0.0)                  | 0 (0.0)          | 18 (7.0)                | 5 (1.8)             |
| Fingolimod                            | 0 (0.0)                  | 0 (0.0)          | 4 (1.5)                 | 1 (0.3)             |
| Ozanimod                              | 0 (0.0)                  | 1 (0.3)          | 0 (0.0)                 | 1 (0.4)             |
| Siponimod                             | 0 (0.0)                  | 1 (0.4)          | 2 (0.6)                 | 2 (0.7)             |
| Teriflunomide                         | 1 (0.4)                  | 6 (2.4)          | 6 (2.5)                 | 0 (0.0)             |

DMT, disease-modifying therapy.

### Table 4. Comparative Switching Outcomes Within 1 Year

| Comparison of proportion of treatment switches (1 year) |              |            |         |
|---|--------------|------------|---------|
|   | Odds ratio   | 95% CI     | P value |
| Cladribine tablets                                      |              | Reference  |         |
| Fingolimod  | 5.66         | 2.90-12.15 | <0.0001 |
| Dimethyl fumarate                                       | 10.81        | 5.66-22.87 | <0.0001 |
| Teriflunomide   | 7.34         | 3.78-15.74 | <0.0001 |
| Comparison of time-to-switch at 1 year                  |              |            |         |
|   | Hazard ratio | 95% CI     | P value |
| Cladribine tablets                                      |              | Reference  |         |
| Fingolimod  | 5.21         | 2.54-10.67 | <0.0001 |
| Dimethyl fumarate                                       | 9.26         | 4.91-17.45 | <0.0001 |
| Teriflunomide   | 6.65         | 3.46-12.79 | <0.0001 |

- Within 2 years, proportions of patients switching treatments were 10.4% on cladribine tablets, 26.9% on fingolimod, 43.5% on dimethyl fumarate, and 34.1% on teriflunomide (Table 5)
- The ORs (95% CI) of 2-year switching vs cladribine tablets were fingolimod 3.25 (2.03-5.32), dimethyl fumarate 6.75 (4.29-10.92), and teriflunomide 4.65 (2.92-7.59) (Table 6)
- HRs (95% CIs) for time to switching at 2 years were statistically significant compared with cladribine tablets (fingolimod 2.99 [1.89-4.71], dimethyl fumarate 5.45 [3.73-7.97], and teriflunomide 4.06 [2.73-6.06], respectively) (Table 6)

### Table 5. Switching Outcomes Within 2 Years

|                                       | Cladribine tablets N (%) | Fingolimod N (%) | Dimethyl fumarate N (%) | Teriflunomide N (%) |
|---------------------------------------|--------------------------|------------------|-------------------------|---------------------|
| Total, N                              | 269                      | 264              | 253                     | 255                 |
| <b>Switch within 2 years</b>          | 28 (10.4)                | 71 (26.9)        | 110 (43.5)              | 87 (34.1)           |
| Total number of patients who switched | 28                       | 71               | 110                     | 87                  |
| <b>Name of DMT switched to</b>        |                          |                  |                         |                     |
| Platform                              | <b>5 (1.9)</b>           | <b>9 (3.6)</b>   | <b>17 (6.8)</b>         | <b>17 (6.7)</b>     |
| Glatiramer                            | 2 (0.7)                  | 7 (2.6)          | 11 (4.4)                | 11 (4.4)            |
| Interferon beta-1a                    | 3 (1.1)                  | 1 (0.3)          | 4 (1.6)                 | 5 (1.9)             |
| Interferon beta-1b                    | 0 (0.0)                  | 0 (0.0)          | 1 (0.2)                 | 1 (0.3)             |
| Peginterferon beta-1b                 | 0 (0.0)                  | 1 (0.5)          | 1 (0.5)                 | 0 (0.0)             |
| Monoclonal antibodies                 | <b>17 (6.3)</b>          | <b>39 (15.0)</b> | <b>43 (17.0)</b>        | <b>41 (16.0)</b>    |
| Alemtuzumab                           | 1 (0.4)                  | 0 (0.0)          | 0 (0.0)                 | 0 (0.0)             |
| Natalizumab                           | 1 (0.4)                  | 4 (1.4)          | 8 (3.3)                 | 10 (4.0)            |
| Ocrelizumab                           | 6 (2.2)                  | 32 (12.0)        | 29 (11.0)               | 23 (9.1)            |
| Oratumumab                            | 9 (3.3)                  | 3 (1.1)          | 6 (2.3)                 | 7 (2.9)             |
| Oral                                  | <b>6 (2.2)</b>           | <b>23 (8.6)</b>  | <b>50 (20.0)</b>        | <b>29 (11.0)</b>    |
| Cladribine tablets                    | 0 (0.0)                  | 2 (0.9)          | 2 (0.7)                 | 5 (1.8)             |
| Dimethyl fumarate                     | 2 (0.7)                  | 8 (2.9)          | 0 (0.0)                 | 8 (3.3)             |
| Diroximel fumarate                    | 1 (0.4)                  | 1 (0.2)          | 32 (13.0)               | 6 (2.5)             |
| Fingolimod                            | 0 (0.0)                  | 0 (0.0)          | 4 (1.7)                 | 1 (0.5)             |
| Ozanimod                              | 0 (0.0)                  | 0 (0.0)          | 0 (0.0)                 | 0 (0.0)             |
| Siponimod                             | 1 (0.4)                  | 1 (0.5)          | 2 (0.6)                 | 2 (0.8)             |
| Teriflunomide                         | 0 (0.0)                  | 0 (0.0)          | 0 (0.0)                 | 0 (0.0)             |

DMT, disease-modifying therapy.

### Table 6. Comparative Switching Outcomes Within 2 Years

| Comparison of proportion of treatment switch (2 years) |              |            |         |
|--|--------------|------------|---------|
|  | Odds ratio   | 95% CI     | P value |
| Cladribine tablets                                     |              | Reference  |         |
| Fingolimod   | 3.25         | 2.03-5.32  | <0.0001 |
| Dimethyl fumarate                                      | 6.75         | 4.29-10.92 | <0.0001 |
| Teriflunomide  | 4.65         | 2.92-7.59  | <0.0001 |
| Comparison of time-to-switch at 2 years                |              |            |         |
|  | Hazard ratio | 95% CI     | P value |
| Cladribine tablets                                     |              | Reference  |         |
| Fingolimod   | 2.99         | 1.89-4.71  | <0.0001 |
| Dimethyl fumarate                                      | 5.45         | 3.73-7.97  | <0.0001 |
| Teriflunomide  | 4.06         | 2.73-6.06  | <0.0001 |

**Acknowledgments:** This study was sponsored by EMD Serono (CrossRef Funder ID: 10.13039/100004755), who reviewed and provided feedback on the abstract. Writing and editorial support for the preparation of this abstract was provided by Natalie Edwards of Health Services Consulting Corporation; funding was provided by the study sponsor. The authors had full control of the abstract and provided their final approval of all content. **Disclosures:** XZ, JC, JA, SK, and GGR: Employees of Genesis Research. Genesis Research received funding from EMD Serono (CrossRef Funder ID: 10.13039/100004755) to conduct the study. XC, EE, ALP, and CL: Employees of EMD Serono, Rockland, MA, USA.